

Figure	Page
12. The number of young striped bass examined by size class, and the average relative abundance (%) of prey in fish stomachs within a size class	104
13. The number of young white perch examined by size class, and the average relative biomass (%) of prey in fish stomachs within a size class	105
14. The number of unidentified <i>Morone</i> species examined by size class, and the average relative biomass (%) of prey in fish stomachs within a size class.....	106
15. The number of young <i>Notropis</i> (minnow) species examined by size class, and the average relative biomass (%) of prey in fish stomachs within a size class.....	107
16. The number of young Atlantic menhaden examined by size class, and the average relative biomass (%) of prey in fish stomachs within a size class	108
17. The number of young Clupids (excluding menhaden) examined by size class, and the average relative biomass (%) of prey in fish stomachs within a size class	109
18. Comparison of Strauss Electivity Index values, for prey item <i>Bosmina</i> , for seven species of finfish at four locations within the lower Roanoke watershed...	110
19. Comparison of Strauss Electivity Index values, for prey item cladocerans (excluding <i>Bosmina</i>), for seven species of finfish at four locations within the lower Roanoke watershed	111
20. Comparison of Strauss Electivity Index values, for prey item rotifers, for seven species of finfish at four locations within the lower Roanoke watershed.....	112
21. Comparison of Strauss Electivity Index values, for prey item copepod nauplii, for seven species of finfish at four locations within the lower Roanoke watershed	113
22. Comparison of Strauss Electivity Index values, for prey item copepodid copepods, for seven species of finfish at four locations within the lower Roanoke watershed	114
23. Comparison of Strauss Electivity Index values, for prey item copepod adults, for seven species of finfish at four locations within the lower Roanoke watershed...	115
24. Comparison of Strauss Electivity Index values, for prey item ostracods, for seven species of finfish at four locations within the lower Roanoke watershed...	116